

Insert Size:- L-122 mm x H-170mm single side printing

Clopidogrel

Cloplat

75 mg Film-Coated Tablet
ANTITHROMBOTIC

FORMULATION :

Each film-coated tablet contains :
Clopidogrel Bisulfate USP
eq. To Clopidogrel 75 mg
Excipients q.s.

PHARMACOKINETICS :

Clopidogrel is rapidly but incompletely absorbed after oral doses; absorption appears to be at least 50%. It is a prodrug and is extensively metabolised in the liver, mainly to the inactive carboxylic acid derivative. The active metabolite appears to be a thiol derivative but has not been identified in plasma. Clopidogrel and the carboxylic acid derivative are highly protein bound. Clopidogrel and its metabolites are excreted in urine and in faeces; about 50% of an oral dose is recovered from the urine and about 46% from the faeces.

INDICATIONS :

Prophylaxis for thromboembolic events; treatment of acute coronary syndromes including unstable angina and non-Q wave myocardial infarction.

DRUG INTERACTION :

Clopidogrel should be used with caution in patients receiving other drugs that increase the risk of bleeding, including anticoagulants, other antiplatelets, and NSAIDs. Clopidogrel may inhibit the cytochrome P450 isoenzyme CYP2C9 and interactions with drugs metabolised by this isoenzyme are theoretically possible; it may also inhibit CYP2B6.

ADVERSE EVENTS :

The incidence of adverse effects, particularly blood dyscrasias, is lower with clopidogrel, although fatalities have been reported. Other adverse effects, reported rarely, include serum sickness, interstitial pneumonitis, erythema multiforme, Stevens-Johnson Syndrome, lichen planus and myalgia.

PRECAUTIONS :

Use with caution in any patient at risk of bleeding, those with active hepatic disease and in patients taking medications that increase the risk of bleeding such as non-steroidal anti-inflammatory agents or aspirin.

Discontinue Clopidogrel 7 days prior to surgery or any event when a normal platelet effect is needed.

DOSAGE AND ADMINISTRATION :

For the prophylaxis of thromboembolic events, the usual dose of clopidogrel is 75 mg once daily. In the management of acute coronary syndromes, including unstable angina and non-Q wave myocardial infarction, clopidogrel is given as a single 300-mg loading dose, followed by 75 mg once daily. Or as prescribed by the physician.

AVAILABILITY :

Clopidogrel (CLOPLAT) 75 mg Film-Coated tablet, packed in alu-alu blister pack by 10's (Box of 30's)

CAUTION :

Foods, Drugs, Devices and Cosmetics Act prohibits dispensing without prescription.

STORE AT TEMPERATURES NOT EXCEEDING 30°C.

Manufactured by :
PSA INTERNATIONAL
(Export Div. of PSA Chemicals
& Pharmaceuticals Pvt. Ltd.)
Vashi, Navi Mumbai, India.

Imported by :
SUHITAS PHARMACEUTICALS INC.
3rd. Floor Centrepoint Bldg.,
Pasonang Tamo, cor. Export Bank Drive,
Makati City, Philippines.

Exclusively distributed by :
 **OEP PHILIPPINES, INC.**
Room No. 606 SEDCO 1 Bldg.,
cor. Rada & Legaspi Sts.,
Legaspi Village, Makati City.